arXiv:1209.2323v2 [nucl-ex] 19 Sep 2012

The event-plane method is obsolete

Matthew Luzum, Jean-Yves Ollitrault
CNRS, URA2306, IPhT, Institut de physique theorique de Saclay, F-91191 Gif-sur-Yvette, France
(Dated: September 20, 2012)

We argue that the traditional event-plane method, which is still widely used to analyze anisotropic flow in ultrarelativistic heavy-ion collisions, should be abandoned because flow fluctuations introduce an uncontrolled bias in the measurement. Instead, one should use an alternative, such as the scalar-product method or cumulant method, which always measures an unambiguous property of the underlying anisotropic flow and therefore eliminates this bias, and does so without any disadvantages. It is known that this correction is important for precision comparisons of traditional v_n measurements requiring better than a few percent accuracy. However, we show that it is absolutely essential for correlations between different harmonics, which can differ from the nominally-measured quantity by as much as 60% to a factor two, depending on the implementation, which is not standard between groups. We also describe how, using the corrected analysis method, the information from different subevents can be combined in order to optimize the precision of analyses.

I. INTRODUCTION

Anisotropic flow v_n, in particular elliptic flow v_2, is one of the most important observables of ultrarelativistic heavy-ion collisions [1–8]. It is therefore important to devise the best possible methods to analyze it. A good measurement must be reproducible; in particular, it must be done in such a way that one can easily compare results from different experiments, using different detectors. For the sake of comparison with theory, it is also important to measure well-defined quantities, corresponding to generic properties of the system that are closely related to an interesting theoretical concept.

In this paper, we argue that the event-plane method [9], which is still used by most of the major experimental collaborations [7, 8, 10–12] does not satisfy any of the above requirements defining a good measurement. When the method was first developed, it was assumed that event-by-event fluctuations in v_n were negligible. However, it is now known that v_n fluctuates significantly within a class of events [13–15]. As a result, an event-plane measurement of v_n yields an ambiguous measure lying somewhere between the event-averaged mean value $\langle v_n\rangle$ and the root-mean-square value $\langle v_n^2\rangle^{1/2}$ [16, 17]. Where exactly depends on the "resolution", which strongly depends on the experimental setup. This means that the same Pb-Pb collisions at the LHC yield a systematically different value of v_n depending on whether they are analyzed by ALICE, CMS or ATLAS, and that comparisons to earlier measurements from lower energy collisions at RHIC are ambiguous. The difference is typically of a few percent for elliptic flow [17], but is larger for higher harmonics like triangular flow v_3 [15], which is solely due to fluctuations.

A slight variant of the event-plane method such as the scalar-product method introduced by the STAR collaboration [18] or equivalently a two-particle cumulant method [19, 20] make for a superior measurement because they consistently yield the rms value of v_n, while introducing no disadvantage compared to the traditional event-plane measurements. Supplemented by higher order cumulant measurements, these make traditional event-plane measurements redundant as well as ambiguous, as they contain no independent information about the underlying flow.

Traditional event-plane analysis methods have also been used to measure higher-order, mixed-harmonic correlations [21–23]. These measurements also suffer from similar biases as traditional event-plane v_n measurements, but in this case they are much larger [24], causing significant problems even for qualitative comparisons. Fortunately, these problems can be repaired with a few simple changes.

In Sec. II, we explain why the traditional event-plane method yields an ambiguous measurement. We show how the ambiguity can be simply removed. We also describe a new systematic way of improving the accuracy of the analysis by combining reference flows from "subevents". In Sec. III, the discussion is extended to the analysis of mixed correlations, where we demonstrate that the proposed corrected procedure is critical.

II. MEASURING v_n

The picture underlying anisotropic flow measurements is that, if the system exhibits strong collective behavior (i.e., "flow"), particles in each event are emitted independently according to an underlying one-particle probability distribution [25]

$$\frac{2\pi}{N}\frac{dN}{d\phi} = 1 + 2\sum_{n=1}^{\infty} v_n \cos n(\phi - \Phi_n), \tag{1}$$

where ϕ is the azimuthal direction of an emitted particle, v_n is the amplitude of anisotropic flow in the nth harmonic, and Φ_n the corresponding reference angle. Further assumptions underlying most analyses are that Φ_n represents a global angle that depends little on pseudorapidity and transverse momentum, and that event-by-event fluctuations of v_n also depend little on pseudorapidity and transverse momentum.

Experiments typically detect anywhere from a few hundred particles to at most a few thousand in a given collision event, while the anisotropy coefficients v_n are typically a few percent or less. A precise reconstruction of the underlying probability distribution is therefore impossible in a single event. Information about the underlying probability distribution can only be extracted from (azimuthally symmetric) correlations between outgoing particles, averaged over a large ensemble of events [26].

Historically, the most common way to access information about the coefficients v_n has been so-called event-plane measurements. In these analyses, v_n is extracted from correlations between particles of interest and a reference detector which is typically separated by a gap in pseudorapidity [4, 27]. In each event, one defines the flow vector $\vec{Q}_n \equiv (|Q_n|\cos(n\Psi_n), |Q_n|\sin(n\Psi_n))$ in harmonic n by [9]

$$\begin{aligned} |Q_n|\cos(n\Psi_n) &= \sum_j w_j \cos(n\phi_j) \\ |Q_n|\sin(n\Psi_n) &= \sum_j w_j \sin(n\phi_j) \end{aligned} \tag{2}$$

where the sum runs over particles seen in the reference detector, with azimuthal angles ϕ_j, and w_j is a weight [1]. Alternatively, if the reference detector is a calorimeter, ϕ_j denotes the azimuthal angle of a segment of the calorimeter, and w_j the energy deposited in the segment.

The idea behind the event-plane method is that the direction of the flow vector Ψ_n provides an estimate of the corresponding angle Φ_n in the underlying probability distribution [29]. Because a finite sample of particles is used, statistical fluctuations cause Ψ_n to differ from Φ_n. The goal is to measure a quantity that is insensitive to these trivial statistical fluctuations and measures an unambiguous property of the underlying probability distribution Eq. (1). In the standard event-plane method, then, the final measurement is defined as

$$v_n\{EP\} \equiv \frac{\langle \cos n(\phi - \Psi_n)\rangle}{Res}, \tag{3}$$

where angular brackets denote an average over particles (with azimuthal angles ϕ) and events. *Res* denotes a resolution correction, which aims to correct for these trivial statistical fluctuations. To do this, one measures correlations between the reference detector in question and one or more additional reference detectors. There are essentially two cases, depending on the detector setup.

1. If there are two identical reference detectors A and B symmetric around midrapidity and separated by a rapidity gap [4] (two-subevent method [9, 29]), then
$$Res \equiv \sqrt{\langle \cos n(\Psi_{nA} - \Psi_{nB})\rangle}. \tag{4}$$
2. If there is no such symmetry, reconstructing the reference flow requires information from a *third* [9] reference detector C, where A, B and C are pairwise separated by pseudorapidity gaps (three-subevent method [7, 11, 12, 22]) The resolution correction for v_n measured with respect to the event plane from detector A is then
$$Res = \sqrt{\frac{\langle \cos n(\Psi_{nA} - \Psi_{nB})\rangle \langle \cos n(\Psi_{nA} - \Psi_{nC})\rangle}{\langle \cos n(\Psi_{nB} - \Psi_{nC})\rangle}}. \tag{5}$$

When the method was developed, it was assumed that dynamical fluctuations of the underlying probability distribution are negligible. As such, the two underlying hypotheses were:

- The numerator of Eq. (3), $\langle \cos n(\phi - \Psi_n)\rangle$, factorizes into $\langle \cos n(\phi - \Phi_n)\rangle \langle \cos n(\Phi_n - \Psi_n)\rangle$.
- The resolution correction *Res* is equal to $\langle \cos n(\Phi_n - \Psi_n)\rangle$.

In this case, it is clear that the resolution correction completely cancels the statistical fluctuations coming from the use of a finite sample of particles in the reference detector, and $v_n\{EP\}$ measures directly $\langle \cos n(\phi - \Phi_n)\rangle$.

However, both hypotheses fail if v_n in the underlying probability distribution fluctuates. The strength of the correlations between ϕ and Φ_n, and between Ψ_n and Φ_n, both increase with v_n, which causes factorization to break down. Similarly, the resolution correction is no longer a simple projection of the measured event plane Ψ_n onto the "true" event plane Φ_n.

In the limit of perfect resolution (i.e., the number of particles used in the reference detector goes to infinity), $\Psi_n = \Phi_n$ and $v_n\{EP\}$ does indeed measure the event-averaged mean v_n from Eq. (1),

$$v_n\{EP\} \to \langle \cos n(\phi - \Phi_n)\rangle = \langle v_n\rangle. \tag{6}$$

In reality, the resolution is not perfect, and the result is usually closer to the low resolution limit [16]. In this limit, the numerator of Eq. (3) is a two-particle correlation, which measures the average value of $\langle v_n \bar{v}_n\rangle$, where $\bar{v}_n$ is the integrated v_n in the reference detector. Similarly, the resolution correction is the square root of a two particle correlation in the reference detector, that is, $\sqrt{\langle \bar{v}_n^2\rangle}$. The event-plane measurement thus yields a root-mean-square value, [2]

$$v_n\{EP\} \to \sqrt{\langle v_n^2\rangle}. \tag{7}$$

In general, the event-plane method yields a result which may lie anywhere between the two limits (6) and

---

[1] The simplest choices are $w_j = 1$ for $n \geq 2$, and $w_j = p_T - \langle p_T^2\rangle/\langle p_T\rangle$ for $n = 1$ [28].

[2] For simplicity we assume here that event-by-event flow fluctuations depend little on rapidity and transverse momentum, so that $v_n/\bar{v}_n$ is the same in every event.



FIG. 1. (Color online) Ratio of the rms v_n, corresponding to the low-resolution limit of Eq. (3) integrated over p_T, to the mean v_n, corresponding to the high-resolution limit, as a function of the number of participant nucleons (binned in 5% centrality intervals), for a Pb-Pb collision at 2.76 TeV per nucleon pair. We model flow fluctuations by assuming $v_n \propto \varepsilon_n$ in each event, where ε_n is the anisotropy of the initial distribution in the corresponding harmonic, defined as in Ref. [30]. The value of ε_n in each event is obtained from the Phobos Monte Carlo Glauber model [31]. For v_3, the ratio is close to $2/\sqrt{\pi} \simeq 1.13$, corresponding to Gaussian fluctuations [32].

(7). The exact measured quantity depends the detector acceptance—even after the resolution correction is applied—so that the measurement is ambiguous.

The difference between the two limits ranges from 6 to 13%, as illustrated in Fig. 1. As the field enters an era of precision physics, comparisons of different experimental analyses [17] and of theory to experiment [33] are significantly complicated by this dependence on analysis details. For example, the large detector acceptance and high multiplicities of recent measurements at the LHC allow for a larger resolution that was available at RHIC. The natural choice (and what has been done) is to take advantage of this extra resolution to minimize the statistical uncertainty of the measurement. The penalty, however, is that comparisons to earlier measurements become ambiguous, and one has difficulty determining from these measurements how much the flow changes, e.g., with collision energy. Theoretical comparisons are hampered for the same reason. Theorists have access to more information than experiments, for example neutral particles in addition to charged. However, if they succumb to the temptation to make use of this extra information to improve statistics, comparisons to measurements become ambiguous and unreliable — even if they painstakingly follow the exact same procedures as the experiment [34–36].

Fortunately, this ambiguity can be removed by a simple change to this procedure, known as the scalar-product method [18]. Although it was originally introduced for unrelated reasons, it turns out to give a well-defined measurement in the presence of flow fluctuations [17], despite differing only trivially from the above analysis. In this method, one simply replaces Eq. (3) with

$$v_n\{SP\} \equiv \frac{\langle |Q_n| \cos n(\phi - \Psi_n)\rangle}{\bar{Q}_n}, \tag{8}$$

where the resolution correction $\bar{Q}_n$ is determined in the same way as the traditional event-plane method except without dividing by the magnitude of the measured flow vector $|\vec{Q}_n|$.

Again there are essentially two cases. For identical and symmetric reference detectors we have

$$\bar{Q}_{nA} \equiv \sqrt{\langle \vec{Q}_{nA} \cdot \vec{Q}_{nB} \rangle}. \tag{9}$$

while more generally one can use

$$\bar{Q}_{nA} \equiv \sqrt{\frac{\langle \vec{Q}_{nA} \cdot \vec{Q}_{nB}\rangle \langle \vec{Q}_{nA} \cdot \vec{Q}_{nC}\rangle}{\langle \vec{Q}_{nB} \cdot \vec{Q}_{nC}\rangle}}. \tag{10}$$

When flow fluctuations are negligible, the scalar product method thus defined is equivalent to the traditional event-plane method. When v_n fluctuates, $\bar{Q}_{nA}$ defined by either (9) or (10) scales like the rms value of $\bar{v}_n$, while the numerator of Eq. (8) scales like $\langle v_n \bar{v}_n \rangle$. Therefore, the scalar-product method always yields the root-mean-square v_n, regardless of the details of the analysis, and makes for a superior measurement.

Equivalently, one can choose to measure a direct two-particle correlation [37], which also always measures this same unambiguous root-mean-square value [13]. It is important to emphasize, however, that even when an event-plane-type analysis is preferred for practical reasons, only a trivial change to the standard analysis is required in order to obtain a well-defined observable.

If the resolution is large enough, higher-order cumulants [19, 20] can be analyzed. These then yield an unambiguous measurement of higher-order, even moments of the distribution of v_n, i.e. $\langle (v_n)^{2k} \rangle$ [13]. Since a standard event-plane method measures something different, one might naively think it useful to perform both analyses in order to obtain the maximum amount of independent information. However, it has been proven [17] that the event-plane method, in addition to being ambiguous, contains no independent information beyond that contained in the first two cumulants, $v_n\{2\}$ and $v_n\{4\}$ (regardless of whether nonflow correlations are present in the system). So there is no reason to use a standard event-plane method in any future analysis.

We end this section with a brief description of how the precision of the scalar-product method can be improved beyond previous implementations. Since at least two reference detectors A and B are needed in order to estimate the reference flow $\bar{Q}_A$, it is tempting to combine the information from both detectors into a single measurement with reduced statistical uncertainty. In the standard event-plane method, this can be done for identical subevents at the expense of additional hypotheses and algebraic complications [9, 38], but there is no known way of recombining non-identical subevents, as in the three-subevent method.

With the scalar-product method, combining reference flows from two reference detectors A and B is straightforward, even if they are not identical. One simply measures v_n independently with respect to A and to B. By linearity of the scalar product, combining both measurements (denoted by $v_{n,A}$ and $v_{n,B}$) amounts to taking a weighted average. We show in Appendix A that the optimal weighting is

$$v_n = \frac{\chi_A^2 v_{n,A} + \chi_B^2 v_{n,B}}{\chi_A^2 + \chi_B^2}, \tag{11}$$

where χ_A is the dimensionless resolution parameter [9, 38, 39]

$$\chi_A \equiv \frac{\bar{Q}_{nA}}{\sqrt{\langle(\vec{Q}_{n,A})^2\rangle - \bar{Q}_{nA}^2}}. \tag{12}$$

When detectors are combined, their resolution parameters χ add in quadrature. Generalization to three reference detectors A, B, C is straightforward.

III. HIGHER-ORDER CORRELATIONS

Fourier harmonics of the azimuthal distribution are now measured up to v_4 [10] at RHIC and up to v_5 [6, 40] and v_6 [7] at LHC. Much additional information is contained in mixed correlations between different harmonics [41–44], which give information about event plane angles Φ_n and will likely play a large part in upcoming measurements. Before rushing into analysis, it is important to define observables for good measurements, which will allow unambiguous comparison between different experiments, and with theory.

An early measurement of this type, performed before the importance of flow fluctuations was noticed, is v_4 with respect to the event-plane of elliptic flow Ψ_2 [21, 22]. The high resolution limit of these measurements is the nominal quantity

$$v_4\{\Psi_2\} \to \langle v_4 \cos 4(\Phi_4 - \Phi_2)\rangle. \tag{13}$$

The actual measured quantity again depends on details such as the event-plane resolution. In addition, there is an ambiguity coming from differences in the implementation of the "resolution correction". The seminal paper [9] proposes two possible implementations, which are equivalent when there are no flow fluctuations, but turn out to be different in the presence of fluctuations. Thus one can obtain either of two expressions for the low-resolution limit of the measurement, depending on the actual implementation. STAR [21] has used Eq. (11) of Ref. [9], which reduces to

$$v_4\{\Psi_2\} \to \frac{\langle v_4 \bar{v}_2^2 \cos 4(\Phi_4 - \Phi_2)\rangle}{\langle \bar{v}_2^2\rangle} \tag{14}$$

in the low-resolution limit, where again $\bar{v}_2$ is integrated over the reference detector. PHENIX instead has used Eq. (14) or Eq. (16) of [9], which yields, in the low-resolution limit,

$$v_4\{\Psi_2\} \to \frac{\langle v_4 \bar{v}_2^2 \cos 4(\Phi_4 - \Phi_2)\rangle}{\sqrt{\langle \bar{v}_2^4\rangle}}. \tag{15}$$

The actual measurement should lie between the high- and low-resolution limits corresponding to the respective implementation. As recently demonstrated in Ref. [33] in full event-by-event hydrodynamic calculations, the dependence on resolution is larger than for the $v_n\{EP\}$ measurements described above. In fact, this should already have been apparent. In addition to $v_4\{\Psi_2\}$, the STAR collaboration also measured a 3-particle correlation $v_4\{3\}$ [21], which corresponds unambiguously to the low-resolution limit of their event-plane measurement. They found that $v_4\{3\}$ is larger than $v_4\{\Psi_2\}$ by 20% for mid central collisions. This difference is much larger than the difference between, e.g., $v_2\{EP\}$ and the low resolution limit $v_2\{SP\}$ in the same experiment (which is roughly 5%).

Further, the difference in implementation, coupled with flow fluctuations, is actually the most likely explanation for the puzzling fact that the STAR measurements are systematically larger than the PHENIX measurements of what was claimed to be the same quantity [22, 24]. The fact that this may not have been fully understood until now underlines the problems arising from ambiguities in experimental analyses combined with the presence of flow fluctuations.

Recently, the ATLAS collaboration has released preliminary measurements of an extensive set of correlations between event planes [23] in Pb-Pb collisions at the LHC, for which they decided to use traditional event-plane methods. To illustrate the large ambiguities present in these measurements, we consider the first of these correlations, which is similar to the earlier measurements of $v_4\{\Psi_2\}$. It is a correlation between the event planes Ψ_4 and Ψ_2, namely, $\langle\cos 4(\Psi_4 - \Psi_2)\rangle$, which is then divided by a resolution factor that is supposed to to unravel the correlation between the underlying reference directions Φ_4 and Φ_2. They thus write:

$$\langle\cos 4(\Phi_4 - \Phi_2)\rangle = \frac{\langle\cos 4(\Psi_4 - \Psi_2)\rangle}{\langle\cos 4(\Psi_4 - \Phi_4)\rangle\langle\cos 4(\Psi_2 - \Phi_2)\rangle}. \tag{16}$$

However, this equation is based on the same hypotheses as the event-plane method, which fail if v_2 and v_4 fluctuate. Specifically, the numerator of the right-hand side does not factorize into $\langle\cos 4(\Psi_4 - \Phi_4)\rangle\langle\cos 4(\Psi_2 - \Phi_2)\rangle\langle\cos 4(\Phi_4 - \Phi_2)\rangle$, and the resolution corrections estimated using standard procedures do not actually correspond to $\langle\cos 4(\Psi_4 - \Phi_4)\rangle$ and $\langle\cos 4(\Psi_2 - \Phi_2)\rangle$.

Specifically, the resolution correction $\langle\cos 4(\Psi_4 - \Phi_4)\rangle$ is defined by Eq. (4) or (5). The resolution correction $\langle\cos 4(\Psi_2 - \Phi_2)\rangle$ corresponds to the convention that PHENX used for $v_4\{\Psi_2\}$, which is defined by a simple generalization of Eq. (4) (or (5)), namely,

$$\langle \cos 4(\Psi_2 - \Phi_2)\rangle \equiv \sqrt{\langle \cos 4(\Psi_{2A} - \Psi_{2B})\rangle}. \tag{17}$$

In the limit where the resolutions on both harmonics are perfect, the quantity measured by ATLAS is indeed the nominal value, corresponding to the left-hand side of Eq. (16):

$$\cos 4(\Psi_4 - \Psi_2)\{EP\} \to \langle \cos 4(\Phi_4 - \Phi_2)\rangle. \tag{18}$$

In the limit of low resolution, however, repeating the same reasoning as in the previous section, what is actually measured is the quantity

$$\cos 4(\Psi_4 - \Psi_2)\{EP\} \to \frac{\langle \bar{v}_4 \bar{v}_2^2 \cos 4(\Phi_4 - \Phi_2)\rangle}{\sqrt{\langle \bar{v}_4^2\rangle \langle \bar{v}_2^4\rangle}}. \tag{19}$$

As before, the measured value is generally between the two limits.[3] Similar expressions can be written for the other correlations in the large set measured by ATLAS. Unlike for measurements of v_n, however, there is no known way to even estimate where between these limits the result lies based only on the reported event-plane resolution.

In order to illustrate the dependence of these measurements on resolution, we use here a simplified calculation, though the results are consistent with more sophisticated calculations [33]. It is known from event-by-event hydrodynamic calculations that p_T-integrated v_2, v_4, and the corresponding reference directions Φ_2 and Φ_4 can be accurately predicted in each event from the initial eccentricities according to [30]

$$\begin{aligned} v_2 e^{2i\Phi_2} &= a\varepsilon_2 e^{2i\Phi_{in,2}} \\ v_4 e^{4i\Phi_4} &= b\varepsilon_4 e^{4i\Phi_{in,4}} + c(\varepsilon_2 e^{2i\Phi_{in,2}})^2, \end{aligned} \tag{20}$$

where a, b, c are real coefficients, and ε_n and $\Phi_{in,n}$ are the initial anisotropy in harmonic n and its reference direction, as defined in Ref. [30].

We thus compute ε_n and $\Phi_{in,n}$ using the PHOBOS Monte Carlo Glauber [31] and take the values of b and c from the viscous hydrodynamic calculation by Teaney and Yan [44], for simplicity taking centrality-independent values of $b = 0.018$ and $c = 0.07$ [4].

The right-hand side of Eqs. (18) and (19) are displayed in Fig. 2 (top, left), as well with their ratio (bottom, left), and similarly for Eqs. (13), (14) and (15) on the right side of the figure. The ratio varies from 1.2 to 2 (in agreement with full event-by-event hydrodynamic calculations of $v_4\{\Psi_2\}$ [33]). Comparing with Fig. 1, one sees that the ambiguity induced by the event-plane method is much worse for mixed correlations that for individual measurements of v_n.



FIG. 2. (Color online) Top, left: low and high resolution limits of the correlation between Φ_2 and Φ_4, defined by Eqs. (19) and (18), versus number of participants. Top, right: low and high resolution limits for integrated $v_4\{\Psi_2\}$, defined by Eqs. (13), (14) and (15). Bottom panels: ratios of the low-resolution to high-resolution results. The nominally-measured values correspond to the high-resolution limit, which can differ from what is actually measured (usually closer to the low-resolution limit) by up to a factor 2. All results are for Au+Au collisions at top RHIC energy.

As a recent illustration of the problems that can arise from this, we point to the correlations calculated in Ref. [45], corresponding to the high resolution limit of the recent preliminary ATLAS measurements — i.e., the quantities that were nominally claimed as measured. They are all clearly smaller in magnitude than the measurements, despite each observable having the correct sign and centrality dependence. Naively, this seems to indicate that the theory needs to be changed somehow to accommodate this difference, but in fact the finite resolution of the experiment could in principle explain the entire discrepancy. Further, while it is straightforward to calculate either the low or the high resolution limit, as done in that work, a calculation that reliably corresponds exactly to the measured quantity adds considerable difficulty. In fact, it is not possible without access to more information about the experimental details than is available.

Fortunately, the ambiguity of the event-plane method can easily be removed by using a straightforward generalization of the scalar-product method. We write the flow vector (2) in complex form, which allows to write compact expressions for mixed correlations:

---

[3] Note that the low-resolution limit (19) lies between −1 and +1 by construction, even though it does not reduce to a simple angular correlation.

[4] Teaney and Yan use cumulants of the initial distribution, which are linear combinations of our moments: with their notation, $b = w_4/\mathcal{C}_4$ and $c = w_{4(22)}/\varepsilon_2^2 + 3(w_4/\mathcal{C}_4)\langle r^2\rangle^2/\langle r^4\rangle$. We take $w_4/\mathcal{C}_4 \simeq 0.018$ and $w_{4(22)}/\varepsilon_2^2 \simeq 0.04$ from their Fig. 2, right, and we estimate $\langle r^2\rangle^2/\langle r^4\rangle \simeq 0.6$ from a Glauber calculation.

$$Q_n = \sum_j w_j e^{in\phi_j}. \tag{21}$$

With this notation, one can unambiguously obtain the right-hand side of Eq. (19) by defining

$$\frac{\langle \bar{v}_4 \bar{v}_2^2 \cos 4(\Phi_4 - \Phi_2)\rangle}{\sqrt{\langle \bar{v}_4^2\rangle\langle \bar{v}_2^4\rangle}} = \frac{\mathrm{Re}\langle (Q_{2A})^2 Q_{4B}^*\rangle}{\sqrt{\langle Q_{4A}Q_{4B}^*\rangle\langle (Q_{2A}Q_{2B}^*)^2\rangle}}, \tag{22}$$

where Re stands for the real part, subscripts A and B refer to two symmetric reference detector separated by a rapidity gap, and Q_{nB}^* denotes the complex conjugate of Q_{nB}. In the same way as for individual measurements of v_n, this method yields a result which is unambiguous and independent of the number of particles selected for analysis.

Generalization of the above discussion to other mixed correlations [23, 41] is straightforward. Specifically, the ATLAS collaboration measures correlations of the type $\langle \cos(c_1\Psi_1 + \cdots + lc_l\Psi_l)\rangle$, which are divided by the product of resolution factors in each harmonic (Eq. (1) of Ref. [23]). Ambiguity is removed by inserting appropriate factors of $|Q_n|$. That is, the numerator should be replaced as

$$\begin{aligned}&\langle \cos(c_1\Psi_1 + \cdots + lc_l\Psi_l)\rangle \\ &\quad\to \langle |Q_1|^{|c_1|}\cdots |Q_l|^{|c_l|}\cos(c_1\Psi_1 + \cdots + lc_l\Psi_l)\rangle,\end{aligned} \tag{23}$$

while the resolution factor in harmonic j, denoted by $Res\{c_j j\Psi_j\}$, should be replaced with a straightforward generalization of Eq. (10):

$$Res\{c_j j\Psi_j\} \to \sqrt{\frac{\langle (Q_{jA}Q_{jB}^*)^{|c_j|}\rangle\langle (Q_{jA}Q_{jC}^*)^{|c_j|}\rangle}{\langle (Q_{jB}Q_{jC}^*)^{|c_j|}\rangle}}. \tag{24}$$

In the case of two identical and symmetric reference detectors, this can again be simplified to use

$$Res\{c_j j\Psi_j\} \to \sqrt{\langle (Q_{jA}Q_{jB}^*)^{|c_j|}\rangle}. \tag{25}$$

With this convention for the resolution factors, the Cauchy-Schwarz theorem ensures that all the two-plane correlations lie between -1 and $+1$, though there is no such restriction for three-plane correlations (such as the correlation between Φ_2, Φ_3 and Φ_5) which could in principle exceed unity. Another option is to scale the numerator as proposed in Ref. [41], which amounts to scaling Eq. (23) by $\bar{Q}_1^{|c_1|}\cdots\bar{Q}_j^{|c_j|}$, where the resolution factors $\bar{Q}_j$ are given by Eq. (9) or (10). All possible choices are related to each other by information contained in standard cumulant measurements of v_n, so it is only necessary to choose one convention. The only important consideration is to choose something that does not contain the ambiguities of a traditional event-plane analysis.

Note that the particular correlation between Φ_2 and Φ_4 studied above can be contaminated by nonflow correlations; Therefore it is important that the flow vectors Q_2 and Q_4 are determined in regions separated by a pseudorapidity gap [23]. Other mixed correlations, such as the correlation between Φ_2 and Φ_3, are not sensitive to nonflow correlations (though the resolution factor in the denominator still requires subevents that are separated in pseudorapidity). The need for gaps in pseudorapidity must be considered on a case-by-case basis [46].

IV. CONCLUSION

We have explained that measurements of anisotropic flow using the event-plane method are ambiguous because the result depends systematically on analysis details. This dependence, which is due to event-by-event flow fluctuations, is typically up to a few percent for v_2 and 10% for v_3 and higher harmonics. For mixed correlations involving event planes from different harmonics, the dependence is much larger — we have demonstrated that the result may vary by as much as a factor 2 depending on analysis details. We have shown that these ambiguities of the event-plane method can be easily repaired with only minor modifications and without any additional complication or drawbacks. We therefore conclude that the traditional event-plane method should be abandoned in all future analyses.

ACKNOWLEDGMENTS

ML is supported by the European Research Council under the Advanced Investigator Grant ERC-AD-267258. We thank Instituto de Fisica, Sao Paulo, where this paper was written, for hospitality, and FAPESP-CNRS for financial support (project 2011/51864-0)

Appendix A: Combining reference flows

In this Appendix, we show that Eq. (11) defines the optimal combination between anisotropic flows measured in detectors A and B using the scalar-product method, in the sense that it minimizes the statistical error.

Eq. (8) can be rewritten as

$$v_n\{SP\} = \frac{\langle \vec{Q}_n \cdot \vec{u}\rangle}{\bar{Q}_n}, \tag{A1}$$

where $\vec{u} \equiv (\cos n\phi, \sin n\phi)$ is the unit vector of the particle in harmonic n. Assuming that $v_n \ll 1$, so that $\vec{u}$ is essentially a random unit vector, and assuming that one measures v_n in a narrow bin which contains at most one particle per event, a straightforward calculation shows that the statistical error on $v_n\{SP\}$ is (for simplicity, we drop the index n from now on)

$$\delta v = \frac{1}{\sqrt{2N}}\frac{\sqrt{\langle\vec{Q}^2\rangle}}{\bar{Q}}, \tag{A2}$$

where N is the number of particles in the bin.

Consider two subevents A and B, separated by a gap in pseudorapidity. We denote by $\vec{Q}_A$ and $\vec{Q}_B$ the corresponding flow vectors in harmonic n, and by $\bar{Q}_A$ and $\bar{Q}_B$ the corresponding resolution corrections. If one measures v_n in a detector that is separated in pseudorapidity from both A and B, the measurement can be done using either A or B as a reference:

$$v_{A,B} \equiv \frac{\langle\vec{Q}_{A,B}\cdot\vec{u}\rangle}{\bar{Q}_{A,B}}. \tag{A3}$$

One can combine linearly the flow vectors and the resolution corrections

$$\begin{aligned}\vec{Q} &\equiv \vec{Q}_A + \lambda\vec{Q}_B\\ \bar{Q} &\equiv \bar{Q}_A + \lambda\bar{Q}_B,\end{aligned} \tag{A4}$$

where λ is arbitrary. The resulting combined measurement is

$$v_{\text{combined}} \equiv \frac{\langle\vec{Q}\cdot\vec{u}\rangle}{\bar{Q}} = \frac{\bar{Q}_A v_A + \lambda\bar{Q}_B v_B}{\bar{Q}_A + \lambda\bar{Q}_B}. \tag{A5}$$

Using Eq. (A2), straightforward algebra shows that the value of λ which minimizes the statistical error on v_{combined} is

$$\lambda = \frac{\bar{Q}_B\langle\vec{Q}_A^2\rangle - \bar{Q}_A\langle\vec{Q}_A\cdot\vec{Q}_B\rangle}{\bar{Q}_A\langle\vec{Q}_B^2\rangle - \bar{Q}_B\langle\vec{Q}_A\cdot\vec{Q}_B\rangle}. \tag{A6}$$

Since $\langle\vec{Q}_A\cdot\vec{Q}_B\rangle = \bar{Q}_A\bar{Q}_B$ by definition, this can be rewritten as

$$\lambda = \frac{\bar{Q}_B(\langle\vec{Q}_A^2\rangle - \bar{Q}_A^2)}{\bar{Q}_A(\langle\vec{Q}_B^2\rangle - \bar{Q}_B^2)} = \frac{\bar{Q}_A\chi_B^2}{\bar{Q}_B\chi_A^2}, \tag{A7}$$

where the resolution parameter χ_A has been defined in Eq. (12). Inserting this value of λ into Eq. (A5), one recovers Eq. (11).

[1] K. H. Ackermann *et al.* [STAR Collaboration], Phys. Rev. Lett. **86**, 402-407 (2001) [nucl-ex/0009011].

[2] B. B. Back *et al.* [PHOBOS Collaboration], Phys. Rev. Lett. **89**, 222301 (2002) [nucl-ex/0205021].

[3] C. Alt *et al.* [NA49 Collaboration], Phys. Rev. C **68**, 034903 (2003) [nucl-ex/0303001].

[4] S. S. Adler *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **91**, 182301 (2003) [nucl-ex/0305013].

[5] K. Aamodt *et al.* [ALICE Collaboration], Phys. Rev. Lett. **105**, 252302 (2010) [arXiv:1011.3914 [nucl-ex]].

[6] S. Chatrchyan *et al.* [CMS Collaboration], Eur. Phys. J. C **72**, 2012 (2012) [arXiv:1201.3158 [nucl-ex]].

[7] G. Aad *et al.* [ATLAS Collaboration], arXiv:1203.3087 [hep-ex].

[8] D. Adamova *et al.* [CERES Collaboration], arXiv:1205.3692 [nucl-ex].

[9] A. M. Poskanzer and S. A. Voloshin, Phys. Rev. C **58**, 1671 (1998) [arXiv:nucl-ex/9805001].

[10] A. Adare *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **107**, 252301 (2011) [arXiv:1105.3928 [nucl-ex]].

[11] S. Chatrchyan *et al.* [CMS Collaboration], arXiv:1204.1409 [nucl-ex].

[12] B. Abelev *et al.* [ALICE Collaboration], arXiv:1205.5761 [nucl-ex].

[13] M. Miller and R. Snellings, nucl-ex/0312008.

[14] B. Alver *et al.* [PHOBOS Collaboration], Phys. Rev. Lett. **98**, 242302 (2007) [nucl-ex/0610037].

[15] B. Alver and G. Roland, Phys. Rev. C **81**, 054905 (2010) [Erratum-ibid. C **82**, 039903 (2010)] [arXiv:1003.0194 [nucl-th]].

[16] B. Alver, B. B. Back, M. D. Baker, M. Ballintijn, D. S. Barton, R. R. Betts, R. Bindel and W. Busza *et al.*, Phys. Rev. C **77**, 014906 (2008) [arXiv:0711.3724 [nucl-ex]].

[17] J. -Y. Ollitrault, A. M. Poskanzer, S. A. Voloshin, Phys. Rev. **C80**, 014904 (2009) [arXiv:0904.2315 [nucl-ex]].

[18] C. Adler *et al.* [STAR Collaboration], Phys. Rev. C **66**, 034904 (2002) [nucl-ex/0206001].

[19] N. Borghini, P. M. Dinh, J. -Y. Ollitrault, Phys. Rev. **C64**, 054901 (2001). [nucl-th/0105040].

[20] A. Bilandzic, R. Snellings, S. Voloshin, Phys. Rev. **C83**, 044913 (2011). [arXiv:1010.0233 [nucl-ex]].

[21] J. Adams *et al.* [STAR Collaboration], Phys. Rev. Lett. **92**, 062301 (2004) [nucl-ex/0310029].

[22] A. Adare *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **105**, 062301 (2010) [arXiv:1003.5586 [nucl-ex]].

[23] J. Jia [ATLAS Collaboration], arXiv:1208.1427 [nucl-ex].

[24] C. Gombeaud and J. -Y. Ollitrault, Phys. Rev. C **81**, 014901 (2010) [arXiv:0907.4664 [nucl-th]].

[25] M. Luzum, J. Phys. G **38**, 124026 (2011) [arXiv:1107.0592 [nucl-th]].

[26] S. Wang, Y. Z. Jiang, Y. M. Liu, D. Keane, D. Beavis, S. Y. Chu, S. Y. Fung and M. Vient *et al.*, Phys. Rev. C **44**, 1091 (1991).

[27] B. Alver *et al.* [PHOBOS Collaboration], Phys. Rev. C **81**, 034915 (2010) [arXiv:1002.0534 [nucl-ex]].

[28] M. Luzum and J. -Y. Ollitrault, Phys. Rev. Lett. **106**, 102301 (2011) [arXiv:1011.6361 [nucl-ex]].

[29] P. Danielewicz and G. Odyniec, Phys. Lett. B **157**, 146 (1985).

[30] F. G. Gardim, F. Grassi, M. Luzum and J. -Y. Ollitrault, Phys. Rev. C **85**, 024908 (2012) [arXiv:1111.6538 [nucl-th]].

[31] B. Alver, M. Baker, C. Loizides and P. Steinberg, arXiv:0805.4411 [nucl-ex].

[32] S. A. Voloshin, A. M. Poskanzer, A. Tang and G. Wang, Phys. Lett. B **659**, 537 (2008) [arXiv:0708.0800 [nucl-th]].

[33] F. G. Gardim, F. Grassi, M. Luzum and J. -Y. Ollitrault, arXiv:1203.2882 [nucl-th].

[34] H. Holopainen, H. Niemi and K. J. Eskola, Phys. Rev. C

83, 034901 (2011). [arXiv:1007.0368 [hep-ph]].
[35] H. Petersen, Phys. Rev. C **84**, 034912 (2011). [arXiv:1105.1766 [nucl-th]].
[36] L. Pang, Q. Wang and X. -N. Wang, Phys. Rev. C **86**, 024911 (2012). [arXiv:1205.5019 [nucl-th]].
[37] K. Aamodt *et al.* [ALICE Collaboration], Phys. Lett. B **708**, 249 (2012) [arXiv:1109.2501 [nucl-ex]].
[38] J. -Y. Ollitrault, nucl-ex/9711003.
[39] J. -Y. Ollitrault, Phys. Rev. D **48**, 1132 (1993) [hep-ph/9303247].
[40] K. Aamodt *et al.* [ALICE Collaboration], Phys. Rev. Lett. **107**, 032301 (2011) [arXiv:1105.3865 [nucl-ex]].
[41] R. S. Bhalerao, M. Luzum and J. -Y. Ollitrault, Phys. Rev. C **84**, 034910 (2011) [arXiv:1104.4740 [nucl-th]].
[42] D. Teaney and L. Yan, Phys. Rev. C **83**, 064904 (2011) [arXiv:1010.1876 [nucl-th]].
[43] R. S. Bhalerao, M. Luzum and J. -Y. Ollitrault, Phys. Rev. C **84**, 054901 (2011) [arXiv:1107.5485 [nucl-th]].
[44] D. Teaney and L. Yan, arXiv:1206.1905 [nucl-th].
[45] Z. Qiu and U. W. Heinz, arXiv:1208.1200 [nucl-th].
[46] R. S. Bhalerao, M. Luzum and J. Y. Ollitrault, J. Phys. G **38**, 124055 (2011) [arXiv:1106.4940 [nucl-ex]].